

Entanglement for rank-2 mixed states

Tobias J. Osborne^{1,*}

¹Centre for Quantum Computer Technology and Department of Physics, University of Queensland 4072, Australia
(Dated: February 1, 2008)

In a recent paper, Rungta *et. al.* [Phys. Rev. A, **64**, 042315, 2001] introduced a measure of mixed-state entanglement called the *I-concurrence* for arbitrary pairs of qudits. We find an exact formula for an entanglement measure closely related to the *I-concurrence*, the *I-tangle*, for all mixed states of two qudits having no more than two nonzero eigenvalues. We use this formula to provide a tight upper bound for the entanglement of formation for rank-2 mixed states of a qubit and a qudit.

PACS numbers: 03.67.-a, 03.65.Ud

Quantum entanglement, as typified by the singlet state $|\Psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$, is a uniquely quantum property of any nonlocal superposition state. Unlike other properties of quantum states, entanglement is singled out by the apparently central role it plays in quantum information processing. Indeed, it is becoming clear that entanglement is a *resource* which can be used to perform quantum communication and computational tasks [1]. To this end, it is of paramount importance that a general theory of entanglement be developed in order to understand how it can and cannot be manipulated. The first step in this program is to develop physically motivated *entanglement measures* to quantify how much entanglement is present in a given state. Many entanglement measures have been proposed, however, for mixed states, they typically involve difficult minimisations.

The development of an efficient formula for the entanglement present in arbitrary mixed states is a crucial goal of quantum information science. Such a formula would lead to great advances in the theory of quantum communication, as well as possibly revealing the correct role entanglement plays in quantum computational speed-up. To date, while there has been a great deal of work on this subject, there is only one easily-computable entanglement measure and it is specific to pairs of qubits [2, 3]. The purpose of this Letter is to add to this situation by providing a formula for one particular bipartite entanglement measure, the *I-tangle*, for mixed states which have at most two nonzero eigenvalues.

The structure of this Letter is as follows. We begin by introducing the concurrence and the tangle for a pair of qubits. We then discuss a generalisation of the concurrence and the tangle for pairs of qudits. The main result of this Letter, a formula for the *I-tangle* of rank-2 mixed states, is then established. We conclude with a corollary of the main result, an upper bound for the entanglement of formation of a rank-2 mixed state of a qubit and a qudit.

Before we discuss the *I-tangle*, we introduce the *concurrence*, a mixed-state entanglement measure for states of a pair of qubits AB [2, 3, 4]. The definition of the concurrence makes use of a specific transformation on

density operators, the *spin-flip* operation, which is defined as follows. Consider an arbitrary mixed state ρ of AB . We define the spin-flip of ρ to be

$$\tilde{\rho} \equiv \text{tr}(\rho^\dagger)I \otimes I - \rho_A^\dagger \otimes I - I \otimes \rho_B^\dagger + \rho^\dagger, \quad (1)$$

where $\rho_A = \text{tr}_B(\rho)$ and $\rho_B = \text{tr}_A(\rho)$ denote the reduced density operators for subsystems A and B respectively. (We have included the trace and hermitian adjoint terms so that the spin-flip operation is defined for *arbitrary* operators acting on AB .) The formula for the spin-flip is applicable to arbitrary bipartite systems, in which case it is called the *universal state inverter* [5].

The spin-flip operation Eq. (1) on a pair of qubits is an example of an antilinear operation. To be more precise, consider a pure state $\rho = |\psi\rangle\langle\psi|$. The spin-flip operation, when applied to this state, is equivalent to the expression $\tilde{\rho} = |\tilde{\psi}\rangle\langle\tilde{\psi}|$, where

$$|\tilde{\psi}\rangle = \sigma^y \otimes \sigma^y (|\psi\rangle)^*, \quad (2)$$

and where σ^y is expressed in the computational basis as $\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, and the complex conjugation is taken in the computational basis. The operation in Eq. (2) is clearly an antilinear operator [11]. The definition of the spin-flip as an antilinear operator extends, via linearity, to all mixed states, $\tilde{\rho} = \overrightarrow{\theta} \rho \overleftarrow{\theta}$, where we have added the arrows above the antilinear operator θ representing the spin-flip to indicate the direction in which it acts. It is worth noting that the description of the spin-flip Eq. (1) in terms of an antilinear operator θ is specific to two qubits.

For pure states $\rho = |\psi\rangle\langle\psi|$, the concurrence C of $|\psi\rangle$ is defined to be $C = |\langle\psi|\tilde{\psi}\rangle| = \sqrt{\langle\psi|\tilde{\rho}|\psi\rangle}$. When the state ρ of the two qubits is mixed, the concurrence C is defined to be a minimum over all pure-state decompositions $\{p_i, |\psi_i\rangle\}$ of ρ :

$$C(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i |\langle\psi_i|\tilde{\psi}_i\rangle|. \quad (3)$$

It is convenient to introduce another entanglement measure closely related to the concurrence, the *tangle* τ [12],

which is also defined as a minimisation over pure-state decompositions:

$$\tau(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i |\langle \psi_i | \tilde{\rho} | \psi_i \rangle|^2. \quad (4)$$

The squared-concurrence satisfies the inequality $C^2 \leq \tau$, which follows from the convexity of $|\langle \psi_i | \tilde{\rho} | \psi_i \rangle|^2 = C^2(|\psi_i\rangle)$. It turns out that the reverse inequality also holds, so that the tangle is equal to the square of the concurrence, $\tau(\rho) = C^2(\rho)$ [13]. (The reverse inequality may be established by noting that there exists a decomposition $\{p_i, |\psi_i\rangle\}$ achieving the minimum in Eq. (3) which has the property that $C(|\psi_i\rangle) = C(|\psi_j\rangle)$ [3]. The inequality follows from substituting this decomposition into the expressions for τ and C^2 .) A simple formula for the concurrence of two qubits is known [3],

$$C(\rho) = \max[0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4], \quad (5)$$

where the λ_i are the square-roots of the singular values, in decreasing order, of the matrix $\rho \tilde{\rho}$.

For a pair of qudits AB we use a variant of the I -concurrence of Rungta *et. al.* [5] to measure the entanglement for mixed states of A and B . The I -concurrence is defined via Eq. (1) [5],

$$C(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \sqrt{\langle \psi_i | \tilde{\rho} | \psi_i \rangle}, \quad (6)$$

where $\rho_i = |\psi_i\rangle\langle\psi_i|$. The entanglement measure we use is a generalisation of the tangle, the I -tangle, defined by

$$\tau(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \langle \psi_i | \tilde{\rho} | \psi_i \rangle. \quad (7)$$

The I -concurrence and the I -tangle are good mixed-state entanglement measures because they satisfy the standard properties usually regarded as essential for a good entanglement measure (see, for example, [6, 7]). The inequality $C^2 \leq \tau$ may be established, by convexity, as for two qubits. Because the equal-entanglement decomposition only exists for pairs of qubits, the I -tangle is not, in general, equal to the square of the I -concurrence. Based on the results of this paper, and the calculations of the I -concurrence for isotropic states [14] we feel that the I -tangle, as defined by a minimisation, is the proper generalisation of the tangle Eq. (4).

The universal state inverter Eq. (1) may be expressed in terms of another formula which will be most useful in the following. Before we write down this formula, however, we need to introduce some definitions. Let $|i\rangle_A$ and $|j\rangle_B$ denote the computational basis states for subsystems A and B , with dimensions d_A and d_B , respectively. For an arbitrary pair $\{|i\rangle_A, |i'\rangle_A\}$, $\{|j\rangle_B, |j'\rangle_B\}$ of the computational basis states of AB we set up the projectors $P_A^{(ii')} = |i\rangle_A\langle i| + |i'\rangle_A\langle i'|$, $P_B^{(jj')} = |j\rangle_B\langle j| + |j'\rangle_B\langle j'|$, and $Q_\alpha = P_A^{(ii')} \otimes P_B^{(jj')}$, where $\alpha = (i, i', j, j')$. Consider

the object $\rho_\alpha = Q_\alpha \rho Q_\alpha$. The operator ρ_α is a positive operator supported on a 2×2 subspace of the Hilbert space of AB spanned by $\{|ij\rangle, |i'j\rangle, |ij'\rangle, |i'j'\rangle\}$. In this way we can think of ρ_α as a subnormalised state of two qubits. The two-qubit spin flip, when applied to ρ_α , gives

$$\tilde{\rho}_\alpha = \vec{\theta}_\alpha \rho_\alpha \overleftarrow{\theta}_\alpha = \sigma^y \otimes \sigma^y (Q_\alpha \rho Q_\alpha)^* \sigma^y \otimes \sigma^y, \quad (8)$$

where $\theta_\alpha = \theta Q_\alpha$ is the antilinear operator representing the spin-flip operation on the 2×2 subspace, and σ^y is naturally defined on the two-dimensional subspaces of A and B respectively. Using these definitions we can write an alternative formula for the universal state inverter,

$$\tilde{\rho} = \sum_\alpha \vec{\theta}_\alpha \rho_\alpha \overleftarrow{\theta}_\alpha, \quad (9)$$

where the sum over α runs over all of the $(d_A(d_A - 1)/2)(d_B(d_B - 1)/2)$ possible choices of pairs of computational basis states. (The reader may verify that Eq. (9) follows from the expression of the universal state inverter as a tensor product of two superoperators of the form $\mathcal{P} \circ \mathcal{T}$. See [5] for further details.)

It is convenient, at this point, to introduce two quantities which will simplify the statement of our main result. Let ρ be a density operator for a pair of qudits having no more than two non-zero eigenvalues. We may write ρ in terms of its eigenvectors,

$$\rho = p|v_1\rangle\langle v_1| + (1-p)|v_2\rangle\langle v_2|. \quad (10)$$

Using these eigenvectors we construct the tensor

$$T_{ijkl} = \text{tr}(\gamma_{ij} \tilde{\gamma}_{kl}), \quad (11)$$

where $\gamma_{ij} = |v_i\rangle\langle v_j|$. We also construct the real symmetric 3×3 matrix M_{ij} whose independent entries are given by

$$\begin{aligned} M_{11} &= \frac{1}{4}T_{1221} + \frac{1}{2}T_{1122} + \frac{1}{4}T_{2112}, \\ M_{12} &= \frac{i}{4}T_{1221} - \frac{i}{4}T_{2112}, \\ M_{13} &= \frac{1}{4}T_{1121} - \frac{1}{4}T_{2122} + \frac{1}{4}T_{1112} - \frac{1}{4}T_{1222}, \\ M_{22} &= -\frac{1}{4}T_{1221} + \frac{1}{2}T_{1122} - \frac{1}{4}T_{2112}, \\ M_{23} &= \frac{i}{4}T_{1121} - \frac{i}{4}T_{1112} + \frac{i}{4}T_{2122} - \frac{i}{4}T_{1222}, \\ M_{33} &= \frac{1}{4}T_{1111} - \frac{1}{2}T_{1122} + \frac{1}{4}T_{2222}. \end{aligned} \quad (12)$$

(The entries of M will be shown to be real in the following.)

We now have all the necessary ingredients required for the statement of our main result.

Theorem 1. *Let ρ be any density operator for a pair AB of qudits, of dimensions d_A and d_B , respectively, having*

no more than two nonzero eigenvalues. The I -tangle τ between A and B is given by the expression

$$\tau(\rho) = \text{tr}(\rho\tilde{\rho}) + 2\lambda_{\min}(1 - \text{tr}(\rho^2)), \quad (13)$$

where $\lambda_{\min}$ is the smallest eigenvalue of the matrix M defined by Eq. (12).

The formula Eq. (13) for the I -tangle is easy to compute for all rank-2 mixed states of a pair of qudits. We have verified this formula by performing the minimisation in Eq. (7) numerically for a large number of random states, and compared the results with the formula Eq. (13). We found agreement to within 10^{-16} .

Proof. The method we use to prove this theorem is similar to that employed by Hill and Wootters [2].

Consider an arbitrary pure state $|\psi\rangle$ which can be written as a linear combination of the two eigenvectors of ρ , $|\psi\rangle = c_1|v_1\rangle + c_2|v_2\rangle$. The I -tangle of $|\psi\rangle$ is given by the expression

$$\tau(\psi) = \langle\psi|\tilde{\sigma}|\psi\rangle = \sum_{\alpha} \langle\psi|\vec{\theta}_{\alpha}|\psi\rangle \langle\psi|\vec{\theta}_{\alpha}^{\dagger}|\psi\rangle, \quad (14)$$

where $\sigma = |\psi\rangle\langle\psi|$, and we have used Eq. (9) to rewrite the spin flip in terms of the antilinear operators θ_{α} . Each of the terms in the sum over α may be written as a trace,

$$\tau(\psi) = \sum_{\alpha} \text{tr}(\omega^* \zeta^{\alpha} \omega \zeta^{\alpha*}), \quad (15)$$

where $\omega_{ij} = c_i c_j^*$ is the density matrix of $|\psi\rangle$ expressed in the $\{|v_1\rangle, |v_2\rangle\}$ basis, and $\zeta_{ij}^{\alpha} = \langle v_i | \vec{\theta}_{\alpha} | v_j \rangle$.

The function on the RHS of Eq. (15) can be extended via linearity to a function f of all 2×2 density matrices ω expressed in terms of the $\{|v_1\rangle, |v_2\rangle\}$ basis, i.e., $f(\omega) = \sum_{\alpha} \text{tr}(\omega^* \zeta^{\alpha} \omega \zeta^{\alpha*})$. The function f has the property that it is equal to the I -tangle for all pure states ψ , $f(\psi) = \tau(\psi)$.

Any 2×2 density operator ω may be expressed in terms of the Pauli matrices via the operator expansion, $\omega = \frac{1}{2}(I + \mathbf{r} \cdot \boldsymbol{\sigma})$, where $r_i = \text{tr}(\omega \sigma^i)$. Substituting this expansion into the expression for f gives the quadratic form

$$f(\omega) = \frac{1}{4} \text{tr}(\Upsilon) + \sum_j r_j L_j + \sum_{j,k} r_j r_k M_{jk}, \quad (16)$$

where $\Upsilon = \sum_{\alpha} \zeta^{\alpha*} \zeta^{\alpha}$,

$$L_j = \text{tr}(\sigma^j \Upsilon), \quad (17)$$

and

$$M_{jk} = \sum_{\alpha} \text{tr}(\sigma^j \zeta^{\alpha*} \sigma^k \zeta^{\alpha*}). \quad (18)$$

Each of the terms in the sum over α in Eq. (18) is a real symmetric matrix, so that M is a real symmetric matrix.

It may be straightforwardly verified that the entries of M are given by Eq. (12).

For the rank-2 density operator ρ , the state-space of the system AB can be considered to be the space of all convex combinations of superpositions of $|v_1\rangle$ and $|v_2\rangle$. If a particular state $|\psi\rangle$ of AB is pure, its corresponding 2×2 density operator in the $\{|v_1\rangle, |v_2\rangle\}$ basis, $\omega = \frac{1}{2}(I + \mathbf{r} \cdot \boldsymbol{\sigma})$, satisfies the condition $|\mathbf{r}|^2 = 1$. In this way, we can think of the entire state-space as the Bloch sphere where the poles are the eigenvectors $|v_1\rangle$ and $|v_2\rangle$. A particular decomposition of ρ may be viewed as the weighted sum of points on the surface of the Bloch sphere, where ρ lies at the centre of mass of the weighted sum. The function f is defined on the entire state-space $|\mathbf{r}|^2 \leq 1$.

When the bipartite system AB is a pair of qubits, there is only one term in the sum Eq. (9), and f reduces to the quadratic form that Hill and Wootters [2] study. In this case, the eigenvalues of M are given by $\pm \frac{1}{2} |\det \zeta|$ and $\frac{1}{4} \text{tr}(\zeta^* \zeta)$, which means that f is convex along two directions and concave along a third. In general, the matrix M will have three positive eigenvalues, so that f is typically convex.

For the purposes of this proof it is essential that a quadratic form g be constructed which agrees with f on pure states ψ which has the additional property that it is convex along two directions and linear along a third. A function g which has these properties may be constructed from f as follows,

$$g(\omega) \equiv f(\omega) - \lambda_{\min}(|\mathbf{r}|^2 - 1), \quad (19)$$

where $\lambda_{\min}$ is the smallest eigenvalue of the matrix M . This function is a quadratic form,

$$g(\omega) = K + \sum_j r_j L_j + \sum_{j,k} r_j r_k N_{jk}, \quad (20)$$

where $N = M - \lambda_{\min} I$, and $K = \frac{1}{4} \text{tr}(\Upsilon) + \lambda_{\min}$. The matrix N which defines the quadratic form g has two positive eigenvalues and one zero eigenvalue so that g is convex along two directions and linear along the third. The quadratic form g has the additional property that it is equal to f for pure states ψ , ($|\mathbf{r}|^2 = 1$).

At this point we recall a theorem due to Uhlmann [8], which concerns functions of density matrices expressed as minimisations over all pure-state decompositions.

Theorem 2. *Let G be a positive real-valued function defined on pure states. The function $\mathcal{G}$, defined for all mixed states ρ , given by*

$$\mathcal{G}(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i G(\psi_i), \quad (21)$$

where the minimisation runs over all pure-state decompositions of ρ , $\{p_i, |\psi_i\rangle\}$, is the largest convex function which agrees with G on pure states $\rho = \psi$.

The I -tangle is expressed as a minimisation over all pure-state decompositions of a density operator, so if we could find the largest convex function which agrees with $f = \tau$ on pure states ψ it is guaranteed to be equal to the I -tangle for all mixed states. We claim that g is precisely this function. To prove this statement we proceed via contradiction. Assume that there is a convex function g' which agrees with f on pure states but which is larger than g . We can write g' in terms of g and a ‘correction’ t

$$g'(\omega) = g(\omega) + t(\omega). \quad (22)$$

In order that g' satisfy the correct boundary conditions $t(\omega)$ must be zero on the set of all pure states (the surface of the Bloch sphere). Consider the coordinate system x_i defined by the three eigenvectors of N where, without loss of generality, we choose the x_3 coordinate to correspond with the direction along which g grows linearly. The condition that g' be convex is equivalent to requiring that the Hessian matrix

$$\mathbf{H} = \frac{\partial^2 g'}{\partial x_i \partial x_j} \quad (23)$$

is positive semidefinite [9]. In order that $\mathbf{H}$ be positive semidefinite, it is necessary (although not sufficient) that the entry $\mathbf{H}_{33}$ satisfies the inequality $\mathbf{H}_{33} \geq 0$ [10]. Because g grows linearly along x_3 , the condition that $\mathbf{H}_{33} \geq 0$ becomes $\frac{\partial^2 t}{\partial x_3^2} \geq 0$, which means that t is convex as a function of x_3 . The boundary conditions, $t(\psi) = 0$, therefore imply that t is everywhere negative. Hence $g' \leq g$, and we have a contradiction. This implies that g is the largest convex function which takes the values $\tau(\psi)$ on the set of all pure states. Therefore, theorem 2 shows that g is equal to the I -tangle.

The expression for the I -tangle τ may be simplified by noting that, for rank-2 ρ , $1 - |\mathbf{r}|^2 = 2(1 - \text{tr}(\rho^2))$. Note, also, that $f(\rho) = \text{tr}(\rho\tilde{\rho})$. Hence we can write $\tau(\rho) = \text{tr}(\rho\tilde{\rho}) + 2\lambda_{\min}(1 - \text{tr}(\rho^2))$. $\square$

The decomposition which achieves the minimum for the I -tangle Eq. (13) consists of two terms. In contradistinction to the case of two qubits, the minimising decomposition will, in general, consist of terms with differing values of τ . This is because the surfaces of constant g will typically be curved, so that the trick of Hill and Wootters cannot be applied (see [2] for the construction of the minimising decomposition when the surfaces of constant g are elliptic cylinders). The construction of the minimising decomposition follows from observing that the function g has the property that it grows linearly in one direction. Consider the line parallel to the eigenvector of N , whose associated eigenvalue is zero, which passes through the density operator ρ . The density operator ρ may be written as a convex sum of the two pure states $|\phi_1\rangle$ and $|\phi_2\rangle$ which lie at either end of the line,

$\rho = q_1|\phi_1\rangle\langle\phi_1| + q_2|\phi_2\rangle\langle\phi_2|$. Because the I -tangle is convex, we obtain the inequality

$$\tau(\rho) \leq q_1\tau(\phi_1) + q_2\tau(\phi_2). \quad (24)$$

However, $\tau = g$ varies linearly in this direction, so that the inequality in Eq. (24) is actually an equality.

When one of the subsystems of the bipartite system AB is a qubit it is possible to obtain a relation between the I -tangle τ and the entanglement of formation $\mathcal{F}$. For pure states ψ of AB the entanglement of formation is given in terms of the I -tangle via

$$\mathcal{F}(\psi) = \mathcal{E}(\tau(\psi)), \quad (25)$$

where $\mathcal{E}(x) = H\left(\frac{1}{2} + \frac{1}{2}\sqrt{1-x}\right)$, and H is the binary entropy function $H(x) = -x \log x - (1-x) \log(1-x)$, where the logarithm is taken to base 2. The function $\mathcal{E}$ is concave and monotone increasing. If we consider the minimising decomposition $\{q_i, |\phi_i\rangle\}$ we constructed in the previous paragraph, we obtain the chain of inequalities

$$\mathcal{F}(\rho) \leq q_1\mathcal{E}(\tau(\phi_1)) + q_2\mathcal{E}(\tau(\phi_2)) \leq \mathcal{E}(\tau(\rho)), \quad (26)$$

where the first inequality follows from the definition of the entanglement of formation, and the second from the fact $\mathcal{E}(g)$ is concave along the line passing through the pure states $|\phi_i\rangle$. This statement is the content of the following corollary:

Corollary 1. *For rank-2 mixed states ρ of a qubit A and a qudit B , the entanglement of formation $\mathcal{F}$ of ρ satisfies the inequality*

$$\mathcal{F}(\rho) \leq H\left(\frac{1}{2} + \frac{1}{2}\sqrt{1-\tau(\rho)}\right). \quad (27)$$

Numerical experiments indicate that the expressions on the LHS and RHS of Eq. (27) usually differ only by about 10^{-4} , so that the inequality is typically very close to an equality; it is not, however, an equality.

We have found a formula for the I -tangle for all states of a pair of qudits having no more than two nonzero eigenvalues. We have also found an upper bound for the entanglement of formation between rank-2 mixed states of a qubit and a qudit. The method we employed to construct these formulae relied crucially on theorem 2. It is tempting to conjecture that the construction of entanglement measures might be simplified by searching for large convex functions which satisfy certain boundary conditions — perhaps as solutions to certain partial differential equations. There are some difficulties with this suggestion, however. We must include in this search space functions which are not analytic at one or more points (the concurrence is not analytic at all points). Nonetheless, perhaps there exists a systematic way to construct good analytic approximations to entanglement measures

expressed as minimisations over pure state decompositions.

I would especially like to thank Carl Caves, Michael Nielsen, and Bill Wootters for their help and for inspirational discussions which led to this work. I would also like to thank Carl Caves for pointing out the substantially simpler proof of Eq. (9). Thanks also to Michael Bremner, Chris Dawson, Jennifer Dodd, Alexei Gilchrist, Duncan Mortimer, and Armin Uhlmann for helpful and stimulating discussions. This work has been funded, in part, by an Australian Postgraduate Award.

* tjo@maths.uq.edu.au

- [1] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996).
- [2] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997).
- [3] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
- [4] W. K. Wootters, Quant. Inf. and Comp. **1**, 27 (2001).
- [5] P. Rungta, V. Bužek, C. M. Caves, and G. J. Milburn, Phys. Rev. A **64**, 042315, 13 (2001).
- [6] G. Vidal, D. Jonathan, and M. A. Nielsen, Phys. Rev. A (3) **62**, 012304 (2000).
- [7] V. Vedral and M. B. Plenio, Phys. Rev. A **57**, 1619 (1998).
- [8] A. Uhlmann, Phys. Rev. A (3) **62**, 032307, 9 (2000), ISSN 1050-2947.
- [9] R. T. Rockafellar, *Convex analysis* (Princeton University Press, Princeton, N.J., 1970).
- [10] R. A. Horn and C. R. Johnson, *Matrix analysis* (Cambridge University Press, Cambridge, 1990), ISBN 0-521-38632-2, corrected reprint of the 1985 original.
- [11] An antilinear operator θ is an operator which satisfies $\theta(c_1|\phi_1\rangle + c_2|\phi_2\rangle) = c_1^*\theta|\phi_1\rangle + c_2^*\theta|\phi_2\rangle$.
- [12] It should be noted that our definition of the tangle differs from the definition used in the literature, where the tangle is simply defined to be *equal* to the squared-concurrence, $\tau = C^2$. As we show, our definition is equivalent to the standard one for two qubits. We use this alternative definition because it naturally generalises to a quantity which is different to the squared-concurrence for pairs of qudits.
- [13] The equality of the tangle and the squared-concurrence was pointed out by Michael Nielsen.
- [14] P. Rungta and C. M. Caves, private communication.